

# PTSD signs and symptoms

Jennifer Sumner, PhD



Potentially life-threatening medical events, like SCAD, can be very stressful and scary. It is common to have a variety of emotional reactions to extremely stressful and traumatic events, including fear, anxiety, sadness, and irritability. For most individuals, these emotional reactions will resolve and get better on their own. However, for some individuals, these symptoms can persist and cause great distress and/or interfere with one's home or work life. Some people can even develop posttraumatic stress disorder (PTSD) after a potentially life-threatening medical event. Although we often think of PTSD as a psychological condition that affects war veterans, PTSD can also be triggered by experiencing an acute medical event.<sup>1</sup> Not everyone who experiences a potentially life-threatening medical event will have PTSD onset—some individuals are more likely to develop it than others. However, it has been found to affect a sizeable number of survivors of these events. For example, it has been estimated that approximately 12% of patients develop symptoms of PTSD after a heart attack.<sup>2</sup>

## There are four types of symptoms of PTSD:

### 1) Reliving the event

These symptoms involve having memories of the traumatic event. They can come back at any time, including during the day and at night (e.g., in the form of nightmares). Sometimes, individuals can even feel like they are going through the event again (flashbacks).

### 2) Avoiding reminders of the event

Individuals may try to avoid situations or people that trigger memories of the traumatic event. They may also try to avoid thinking or talking about the event.

### 3) Negative changes in beliefs and feelings

These symptoms involve alterations in how individuals think and feel about themselves, others, and the world. For example, people may no longer experience loving feelings toward others or think that the world is a totally dangerous place.

### 4) Feeling keyed up

Individuals may be keyed up and on alert for danger after a traumatic event. They may startle easily (e.g., jump if there is a loud sound), have difficulty sleeping, or have trouble concentrating.

# PTSD signs and symptoms

Jennifer Sumner, PhD



If you experience symptoms like these after a traumatic event for more than four weeks, find that these symptoms bother you greatly, and/or disrupt your home or work life, it is recommended that you seek professional help from a doctor or counselor. Some resources for finding treatment providers in your area include:

Association for Behavioral and Cognitive Therapies “Find a CBT Therapist”

<http://www.findcbt.org/xFAT/>

International Society for Traumatic Stress Studies Clinician Directory

<http://www.istss.org/find-a-clinician.aspx>

**Jennifer Sumner, PhD**, is an Instructor at the Center for Behavioral Cardiovascular Health at Columbia University Medical Center and a Visiting Scientist at the Harvard T.H. Chan School of Public Health. She is a licensed clinical psychologist whose work focuses on the mechanisms by which trauma exposure contributes to negative outcomes for emotional and physical health. She takes a life-course perspective on trauma, studying mechanistic processes in children, adolescents, and adults. Her recent work has examined the pathways by which trauma exposure and posttraumatic stress disorder (PTSD) contribute to cardiovascular disease, particularly in women. She is the recipient of a Career Development Award from the National Heart Lung and Blood Institute to pursue this work. Her research provides evidence that PTSD is not solely a mental health problem but also increases risk of chronic disease, and it demonstrates the need for integrated mental and physical health care. She received her BA from Pomona College and her PhD from Northwestern University.

## References

- 1 Edmondson D, Cohen BE. Posttraumatic stress disorder and cardiovascular disease. *Prog Cardiovasc Dis.* 2013;55:548-556.
- 2 Edmondson D, Richardson S, Falzon L, Davidson KW, Mills MA, Neria Y. Posttraumatic stress disorder prevalence and risk of recurrence in acute coronary syndrome patients: a meta-analytic review. *PLoS One.* 2012;7:e38915.